

# Lumpings of Markov chains and entropy rate loss

Bernhard C. Geiger<sup>\*†</sup> & Christoph Temmel<sup>‡§</sup>

20130115

## Abstract

A lumping of a Markov chain is a coordinate-wise projection of the chain. We characterise the entropy rate loss of a lumping of an aperiodic and irreducible Markov chain on a finite state space in two ways. First, by the random growth rate of the number of realisable preimages of a finite-length trajectory of the lumped chain. Second, by the possibility to reconstruct original trajectories from their lumped images. Both are purely combinatorial criteria, depending only on the transition graph of the Markov chain and the lumping function. We state sufficient conditions on a non-positive transition matrix and a lumping to preserve the entropy rate. In the sparse setting, we give sufficient conditions on the lumping to both preserve the entropy rate and result in  $k$ -th order homogeneous Markov chain. Every non-trivial lumping of a Markov chain with positive transition matrix incurs an entropy rate loss.

Keywords: lumping, entropy rate loss, functional hidden Markov model, lumpability

MSC 2010: 60J10 (primary), 60G17, 94A17, 60G10, 65C40

## Contents

|          |                                                                          |          |
|----------|--------------------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                                      | <b>3</b> |
| <b>2</b> | <b>Main results</b>                                                      | <b>3</b> |
| 2.1      | Preliminaries . . . . .                                                  | 3        |
| 2.2      | Setting . . . . .                                                        | 4        |
| 2.3      | The characterisation of entropy rate loss . . . . .                      | 5        |
| 2.4      | Sufficient conditions for entropy rate preservation and $k$ -lumpability | 6        |
| 2.5      | Further discussion . . . . .                                             | 7        |

---

<sup>\*</sup>4420 Signal Processing and Speech Communication Laboratory, Graz University of Technology, Inffeldgasse 16c, 8010 Graz, Austria

<sup>†</sup>Email: geiger@ieee.org

<sup>‡</sup>5030 Institute for Mathematical Structure Theory, Graz University of Technology, Steyrergasse 30/III, 8010 Graz, Austria

<sup>§</sup>Email: temmel@math.tugraz.at

|          |                                                            |           |
|----------|------------------------------------------------------------|-----------|
| <b>3</b> | <b>Proof of the characterisation</b>                       | <b>10</b> |
| 3.1      | The preservation case . . . . .                            | 10        |
| 3.2      | Non-overlapping traversal instants . . . . .               | 11        |
| 3.3      | Conditional Markov property . . . . .                      | 13        |
| 3.4      | The loss case . . . . .                                    | 15        |
| <b>4</b> | <b>Proofs of the sufficient conditions</b>                 | <b>16</b> |
| 4.1      | Single entry implies entropy rate preservation . . . . .   | 16        |
| 4.2      | Single forward property implies $k$ -lumpability . . . . . | 17        |

## List of Figures

|   |                                                                                               |   |
|---|-----------------------------------------------------------------------------------------------|---|
| 1 | Illustration of $\mathcal{K}$ . . . . .                                                       | 5 |
| 2 | SE is neither necessary for entropy rate preservation nor for weak $k$ -lumpability . . . . . | 7 |
| 3 | SFS( $k$ ) is not necessary for entropy rate preservation and strong lumpability . . . . .    | 8 |
| 4 | SE does neither imply SFS( $k$ ) nor $k$ -lump . . . . .                                      | 8 |

## 1 Introduction

The *entropy rate* of a stationary stochastic process is the average number of bits per time unit needed to encode the process. A *lumping* of a (stationary) Markov chain is a coordinate-wise projection of the chain by a *lumping function*. The resulting (stationary) *lumped stochastic process* is also called a *functional hidden Markov model* [EM02]. One can transform an arbitrary hidden Markov model on finite state and observation spaces into this setting [EM02, Section IV.E]. In general, the lumped process loses the Markov property [GL05] and has a lower entropy rate than the original Markov chain due to the aggregation of states [Pin64, WA60].

The present paper investigates the structure of entropy rate preserving lumpings of stationary Markov chains over a finite state space. The central result characterises the entropy rate preserving case in two ways. First, by a structural condition on the transition graph associated with the Markov chain. Second, by the growth of the number of realisable preimages of a realisation of the lumped process. We document a strong dichotomy between the preservation and loss case: uniform finite bounds on the lost entropy and the number of realisable preimages in the former and a linearly growing entropy loss and an almost-surely exponentially growing number of realisable preimages in the latter.

In particular, a positive transition matrix always implies an entropy rate loss for a non-identity lumping. We state sufficient conditions on a lumping of a Markov chain with a sparse transition graph to preserve the entropy rate. The representation of each finite-state stationary stochastic process as a lumping of Markov chain on an at most countable state space by Carlyle [Car67] fulfils this condition.

More refined sufficient conditions additionally yield higher-order Markov behaviour of the lumped process. This behaviour is highly desirable from a simulation point of view. The sufficient conditions for sparse transition matrices complement Gurvits & Ledoux's [GL05] result that lumpings of positive transition matrices having higher-order Markov behaviour are nowhere dense.

## 2 Main results

### 2.1 Preliminaries

We write  $[n, m[ := \{k \in \mathbb{Z} : n \leq k < m\}$  and all obvious variations thereof. In particular, we abbreviate  $[n] := [1, n]$ .

We briefly review the information-theoretic basics from Cover & Thomas [CT06]. Let  $\text{ld}$  denote the binary logarithm. By continuous extension we assume  $0 \text{ ld } 0 = 0$ . The *Shannon entropy* [CT06, (2.1)] of a rv  $Z$  taking values in a finite set  $\mathcal{Z}$  is

$$H(Z) := - \sum_{z \in \mathcal{Z}} \mathbb{P}(Z = z) \text{ld } \mathbb{P}(Z = z). \quad (1a)$$

The *conditional entropy* [CT06, (2.10)] of  $Z$  given  $W$  is defined by

$$H(Z|W) := \sum_{w \in \mathcal{W}} \mathbb{P}(W = w) H(Z|W = w). \quad (1b)$$

Successive conditioning reduces entropy [CT06, Theorem 2.6.5]:

$$H(Z) \geq H(Z|W_1) \geq H(Z|W_1, W_2). \quad (1c)$$

For a stationary stochastic process  $Z := (Z_n)_{n \in \mathbb{Z}}$  on a finite state space  $\mathcal{Z}$ , the *entropy rate* [CT06, Theorem 4.2.1, pp. 75] is

$$\overline{H}(Z) := \lim_{n \rightarrow \infty} \frac{1}{n} H(Z_{[n]}) = \lim_{n \rightarrow \infty} H(Z_n | Z_{[n-1]}). \quad (1d)$$

The left limit in (1d) is the limit of the normalised *block entropy*  $H(Z_{[n]})$ . By stationarity and (1c) the  $H(Z_n | Z_{[n-1]})$  in the right limit are monotonically decreasing.

## 2.2 Setting

This section describes the setting of our work. Let  $X := (X_n)_{n \in \mathbb{Z}}$  be an irreducible, aperiodic, homogeneous Markov chain on the finite state space  $\mathcal{X}$ . It has transition matrix  $P$  with invariant probability measure  $\mu$ . We assume that  $X$  is stationary, that is  $X_0 \sim \mu$ . The *lumping function*  $g$  is  $\mathcal{X} \rightarrow \mathcal{Y}$  and surjective. We assume  $g$  to be non-trivial, that is  $2 \leq |\mathcal{Y}| < |\mathcal{X}|$ . Without loss of generality, we extend  $g$  to  $\mathcal{X}^n \rightarrow \mathcal{Y}^n$ , for arbitrary  $n \in \mathbb{N}$ . The *lumped process* of  $X$  under  $g$  is the stochastic process  $Y := (Y_n)_{n \in \mathbb{Z}}$  defined by  $Y_n := g(X_n)$ . We refer to this setup also as the *lumping*  $(P, g)$ .

The lumping induces a *conditional entropy rate* [GK11, WA60], which characterises the average information loss per time unit:

$$\overline{H}(X|Y) := \lim_{n \rightarrow \infty} \frac{1}{n} H(X_{[n]} | Y_{[n]}) = \overline{H}(X) - \overline{H}(Y). \quad (2)$$

Our main question is whether  $\overline{H}(X|Y)$  is positive or zero. We speak of *entropy rate loss* or *entropy rate preservation* respectively. Note that  $\overline{H}(X|Y) = 0$  does not imply that the original process can be reconstructed from the lumped process (see figure 3 for an example).

Recall that the *transition graph*  $G$  of the Markov chain  $X$  is the directed graph with vertex set  $\mathcal{X}$  and an edge  $(x, x')$ , iff  $P_{(x, x')} > 0$ . Our results partly depend only on the structure of  $G$ , that is, whether an entry of  $P$  is positive or zero. In particular, if we want to investigate the number of realisable preimages of a given finite-length lumped trajectory, then this number only depends on the structure of  $G$ . Informally, multiple realisable preimages are intimately connected with a loss of entropy when passing through the lumped trajectory of finite length. To describe up to which length lumped trajectories have unique realisable preimages, given knowledge of the original endpoints we introduce a key quantity of the lumping  $(P, g)$ :

$$\mathcal{K} := \inf \left\{ n \in \mathbb{N} \mid \begin{array}{l} \exists \tilde{x}, \hat{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}^n : \exists \mathbf{x}', \mathbf{x}'' \in g^{-1}(\mathbf{y}) : \\ \text{st both } \begin{cases} \mathbb{P}(X_0 = \tilde{x}, X_{[n]} = \mathbf{x}', X_{n+1} = \hat{x}) > 0 \\ \mathbb{P}(X_0 = \tilde{x}, X_{[n]} = \mathbf{x}'', X_{n+1} = \hat{x}) > 0 \end{cases} \end{array} \right\}. \quad (3)$$




Figure 1: (Colour online) An example of  $\mathcal{K} = 3$  with two realisable length-5 trajectories  $(\tilde{x}, x'_1, x'_2, x'_3, \hat{x})$  and  $(\tilde{x}, x''_1, x''_2, x''_3, \hat{x})$  with the same lumped image  $(g(\tilde{x}), y_1, y_2, y_3, g(\hat{x}))$ . The lumped images  $\{g(\tilde{x}), y_1, y_2, y_3, g(\hat{x})\}$  do not have to be distinct and may even all be the same.

Thus  $\mathcal{K} \in \mathbb{N} \cup \{\infty\}$ . Figure 1 on page 5 gives an example of  $\mathcal{K} = 3$ .

### 2.3 The characterisation of entropy rate loss

This section presents the characterisation of the entropy rate loss of a lumping in terms of  $\mathcal{K}$  and the preimage count. The *preimage count of length  $n$*  of the lumping  $(P, g)$  is the random variable counting the realisable preimages of the lumped trajectory of length  $n$ . Using Iverson brackets, this is

$$T_n := \sum_{\mathbf{x} \in g^{-1}(Y_{[n]})} [\mathbb{P}(X_{[n]} = \mathbf{x}) > 0]. \quad (4)$$

Our main result is

**Theorem 1.**

$$\overline{H}(X|Y) > 0 \Leftrightarrow \mathcal{K} < \infty \Leftrightarrow \exists C > 1 : \quad \mathbb{P}(\liminf_{n \rightarrow \infty} \sqrt[n]{T_n} \geq C) = 1, \quad (5a)$$

$$\overline{H}(X|Y) = 0 \Leftrightarrow \mathcal{K} = \infty \Leftrightarrow \exists C < \infty : \quad \mathbb{P}(\sup_{n \rightarrow \infty} T_n \leq C) = 1. \quad (5b)$$

The proofs of all statements in this section are in section 3. The constants in theorem 1 are explicit functions of  $(P, g)$ ; see (37) for (5a) and (15) for (5b). Likewise, an explicit lower bound for the entropy rate loss in case (5a) is stated in (34). Two further results, refining the dichotomy in theorem 1, are uniform bounds on the conditional block entropies contingent on  $\mathcal{K}$  and an upper bound on  $\mathcal{K}$  in the loss case.

**Proposition 2.** *We have*

$$\forall n \leq \mathcal{K} + 1 : \quad H(X_{[n]}|Y_{[n]}) \leq 2 \log(|\mathcal{X}| - |\mathcal{Y}| + 1). \quad (6)$$

**Proposition 3.** *In case (5a), we have*

$$\mathcal{K} \leq \sum_{y \in \mathcal{Y}} |g^{-1}(y)| (|g^{-1}(y)| - 1). \quad (7)$$

Theorem 1 and the following propositions reveal a following dichotomy in behaviour of the entropy of the lumping. If  $\mathcal{K}$  is infinite, then arbitrarily long

trajectories of  $X$  can be reconstructed from the lumped trajectory and knowledge of the endpoints. Therefore, the only entropy loss occurs at those endpoints and is finite. This yields uniform finite bounds on the conditional block entropies and the preimage count. On the other hand, if  $\mathcal{K}$  is finite (see figure 1), then at least two realisable length- $(\mathcal{K}+2)$  trajectories of  $X$  with the same lumped image bifurcate and join. This situation leads to a finite entropy loss. Reasoning based on the ergodic theorem ensures that this situation occurs linearly often in the block length, thus leading to a linear growth of the conditional block entropy. This translates into an entropy rate loss.

Another interesting and direct corollary is

**Corollary 4.** *If  $P$  is positive, i.e., all its entries are positive, then  $\overline{H}(X|Y) > 0$ .*

*Proof.* If  $P$  is positive, then  $\mathcal{K} = 1$ .  $\square$

Thus, the search for entropy rate preserving lumpings must be in the space of sufficiently sparse transition matrices  $P$ .

## 2.4 Sufficient conditions for entropy rate preservation and $k$ -lumpability

We present easy-to-check sufficient conditions for the preservation of the entropy rate. Their proofs are in section 4. Our conditions depend only on the transition graph  $G$  and the lumping function  $g$ . The conditions are only stated in a “forward form”; applying the conditions to the time-reversed setting yields a set of mirrored sufficient conditions, which we omit.

Our first sufficient condition is

**Definition 5.** *A lumping  $(P, g)$  is single entry (short: SE), iff*

$$\forall y \in \mathcal{Y}, x \in \mathcal{X} : \exists! x' := x'(x, y) \in g^{-1}(y) : \forall x'' \in g^{-1}(y) \setminus \{x'\} : \mathbb{P}(X_1 = x'' | X_0 = x) = P_{(x, x'')} = 0. \quad (8)$$

The class of single entry lumpings is entropy rate preserving:

**Proposition 6.** *If  $(P, g)$  is SE, then  $\overline{H}(X|Y) = 0$ .*

Figure 2 on page 7 shows that SE is not necessary for entropy rate preservation.

The case of the lumped process retaining the Markov property is desirable from a computational and modelling point of view. However, in general, the lumped process  $Y$  does not possess the Markov property [KS76, GL05]. Its distribution is more complex, despite living on a smaller set. Thus one may hope that the lumped process belongs to the larger and still desirable class of higher-order Markov chains.

**Definition 7.** *A stochastic process  $Z := (Z_n)_{n \in \mathbb{N}}$  is a  $k$ -th order homogeneous Markov chain (short:  $Z$  is HMC( $k$ )) iff*

$$\forall n \in \mathbb{Z}, k \leq m \in \mathbb{N}, z_n \in \mathcal{Z}, z_{[n-m, n-1]} \in \mathcal{Z}^m : \mathbb{P}(Z_n = z_n | Z_{[n-m, n]} = z_{[n-m, n]}) = \mathbb{P}(Z_n = z_n | Z_{[n-k, n]} = z_{[n-k, n]}). \quad (9)$$




Figure 2: (Colour online) The transition graph of a Markov chain with the lumping represented by red boxes. The lumping is neither SE (violated by transitions from  $a$  into  $B$ ) nor its mirror condition for the time-reversed process (violated by transitions from  $D$  to  $f$ ). On the other hand, the existence of the uniquely represented states  $C_1$  and  $C_2$  allows to distinguish between the trajectories  $(a, b_1, c_1, d_1, f)$  and  $(a, b_2, c_2, d_2, f)$ . Therefore the lumping preserves the entropy rate. Furthermore, this lumping is weakly 1-lumpable. Hence it shows that SE is neither necessary for entropy rate preservation nor for weak  $k$ -lumpability. This also applies to SFS( $k$ ) (see definition 9), which is a subclass of SE.

**Definition 8** (Extension of [KS76, Def. 6.3.1]). A lumping  $(P, g)$  is weakly  $k$ -lumpable (short:  $X$  is  $k$ -lump), iff  $Y$  is HMC( $k$ ). If this holds for each distribution of  $X_0$ , then we call  $(P, g)$  strongly  $k$ -lumpable.

**Definition 9.** For  $k \geq 2$ , a lumping  $(P, g)$  has the single forward  $k$ -sequence property (short: SFS( $k$ )), iff

$$\begin{aligned} \forall \mathbf{y} \in \mathcal{Y}^{k-1}, y \in \mathcal{Y} : \exists! \mathbf{x}' := \mathbf{x}'(\mathbf{y}, y) \in g^{-1}(\mathbf{y}) : \\ \forall x \in g^{-1}(y), \mathbf{x} \in g^{-1}(\mathbf{y}) \setminus \{\mathbf{x}'\} : \\ \mathbb{P}(X_{[k-1]} = \mathbf{x} | Y_{[k-1]} = \mathbf{y}, X_0 = x) = 0, \end{aligned} \quad (10)$$

i.e., iff observing a length- $k$  lumped trajectory uniquely determines the last  $(k-1)$  elements of its preimage.

The single forward  $k$ -sequence property implies entropy rate preservation and  $k$ -lumpability:

**Proposition 10.** If  $(P, g)$  is SFS( $k$ ), then it is  $k$ -lump and SE.

Figure 2 on page 7 shows that SFS( $k$ ) is not necessary for weak 1-lumpability and entropy rate preservation. Figure 3 on page 8 demonstrates that SFS(2) is neither necessary for SE nor for strong 1-lumpability. Finally, figure 4 on page 8 shows that SE does neither imply SFS( $k$ ) nor  $k$ -lump, for every  $k$ .

## 2.5 Further discussion

Functions of Markov chains have been considered in the literature for a long time: In particular, whether the function of a Markov chain possesses the Markov




Figure 3: (Colour online) The transition graph of a Markov chain with the lumping represented by red boxes. The lumping is **SE** and thus preserves the entropy rate. Furthermore, it is strongly 1-lumpable and thus  $H(Y_1|X_0) = H(Y_1|Y_0)$  (see proposition 15). However, observing an arbitrarily long trajectory of the lumped process does not determine the current preimage state. Whence  $(P, g)$  is not **SFS**( $k$ ), for every  $k$ . Therefore, **SFS**( $k$ ) is not necessary for entropy rate preservation and strong lumpability.




Figure 4: (Colour online) The transition graph of a Markov chain with the lumping represented by red boxes. The lumping is **SE**. The loops at  $b_1$  and  $b_2$  imply that the lumped process is not **HMC**( $k$ ), for every  $k$ . This is easily seen by the inability to differentiate between  $n$  consecutive  $b_1$ 's and  $n$  consecutive  $b_2$ 's. When starting in  $B$  and as long as  $P_{(b_1,a)} \neq P_{(b_2,a)}$ , this long sequence of  $B$ s prevents determining the probability of entering  $A$ . Thus this is neither **SFS**( $k$ ) nor  $k$ -lump, for each  $k$ .

property or not has been investigated in, among others, [BR58, RP81]. Kemeny & Snell [KS76] coined the term *lumpability* for this case. Higher-order lumpability, as we use it in this work, has been analysed by Gurvits & Ledoux [GL05], and they showed that the class of Markov chains being lumpable is nowhere dense.

A related problem is the *identification problem*, initially posed by Blackwell & Koopmanns [BK57] (for a recent treatment see [And99]): given a stationary process on a finite state space, is it representable by a lumping of a Markov chain? There are two results from research into this topic having a connection to the present work.

First, Carlyle [Car67] showed that every stationary stochastic process on a finite state space can be represented as a lumping of a Markov chain on an at most countable state space. If the representation involves a Markov chain on a finite state space, then it is SE proposition 6 guarantees entropy rate preservation of the representation.

Second, Gilbert [Gil59] showed that the distribution of a lumping of finite-state Markov chain is uniquely determined by the distribution of  $m$  consecutive samples, where  $m$  depends on the cardinalities of the input and output alphabet. This does not contradict the nowhere dense result of Gurvits & Ledoux, however, since the construction of the process distribution is different from a product of conditional distributions (as it is in the case of lumpability).

Moreover, the nowhere dense property does not prevent our results from being practically relevant. In particular, our sufficient condition holds for complete lower-dimensional subspaces of the space of Markov transition matrices. In other words, if the transition matrix is sufficiently sparse, one can hope that the lumping satisfies some of our sufficient conditions. More generally, one can hope that for a given Markov model there exists a lumping function with a desired output alphabet such that the resulting lumping satisfies our sufficient conditions. Sparse transition matrices are used, e.g., in  $n$ -gram models in automatic speech recognition [BdM<sup>+</sup>92, Table 1], chemical reaction networks [HMMW10, HRSS10, Wil11] and link prediction and path analysis [Sar00].

Although we treat only the stationary case, we expect that our results are generalisable to the case where the Markov chain is time-homogeneous, irreducible and aperiodic, but does not start in equilibrium, i.e., where the initial distribution does not coincide with the invariant measure. Entropy rates exist for a larger class of non-stationary stochastic processes, namely those *stationary in the asymptotic mean* [KR81, Gra90] (short: AMS). Finite state Markov chains and lumpings thereof are AMS [KR81]. A more difficult question would be if aperiodicity could be dropped.

### 3 Proof of the characterisation

*Proof of theorem 1.* The proof of the preservation case (5b) of theorem 1 and of proposition 2 are in section 3.1. The proof of the loss case (5a) of theorem 1 and of proposition 3 are in section 3.4. Sections 3.2 and 3.3 contain technical groundwork and results about Markov chains needed in the proof of the loss case in section 3.4.

The statement in (5) follows from the mutually exhaustive implications

$$\mathcal{K} < \infty \Rightarrow \overline{H}(X|Y) > 0, \quad (11a)$$

$$\mathcal{K} = \infty \Rightarrow \overline{H}(X|Y) = 0 \quad (11b)$$

and

$$\mathcal{K} < \infty \Rightarrow \exists C > 0 : \mathbb{P}(\liminf_{n \rightarrow \infty} \sqrt[n]{T_n} \geq C) = 1, \quad (12a)$$

$$\mathcal{K} = \infty \Rightarrow \exists C > 0 : \mathbb{P}(\sup_{n \rightarrow \infty} T_n \leq C) = 1. \quad (12b)$$

The proofs of implications (11b) and (12b) are in section 3.1 and the proofs of implications (11a) and (12a) are in section 3.4.  $\square$

#### 3.1 The preservation case

The definition of  $\mathcal{K}$  in (3) implies that lumped trajectories of length less than  $\mathcal{K}$  have a unique preimage contingent on the endpoints, i.e., if  $n < \mathcal{K}$ , then  $\forall \tilde{x}, \hat{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}^n$ :

$$\begin{aligned} \mathbb{P}(X_0 = \tilde{x}, Y_{[n]} = \mathbf{y}, X_{n+1} = \hat{x}) &> 0 \\ \Rightarrow \exists! \mathbf{x} \in \mathcal{X}^n : \mathbb{P}(X_{[n]} = \mathbf{x} | X_0 = \tilde{x}, Y_{[n]} = \mathbf{y}, X_{n+1} = \hat{x}) &= 1. \end{aligned} \quad (13)$$

*Proof of proposition 2.* Recall that we assume  $n \leq \mathcal{K} + 1$ . The unique preimage (13) implies that the conditional entropy of the interior of a block, given its lumped image and the states at its ends, is zero:

$$\begin{aligned} &H(X_{[2,n-1]} | X_1, X_n, Y_{[n]}) \\ &= \sum_{\substack{\mathbf{y} \in \mathcal{Y}^n \\ \tilde{x}, \hat{x} \in \mathcal{X}}} \mathbb{P}(X_1 = \tilde{x}, X_n = \hat{x}, Y_{[n]} = \mathbf{y}) \underbrace{H(X_{[2,n-1]} | X_1 = \tilde{x}, X_n = \hat{x}, Y_{[n]} = \mathbf{y})}_{=0 \text{ by (13)}} \\ &= 0. \end{aligned} \quad (14)$$

We apply the chain rule of entropy (cf. [CT06, pp. 22]) to decompose the conditional block entropy into its interior and its boundary. The interior vanishes by (14) and the entropy at the endpoints is maximal for the uniform distribution:

$$\begin{aligned} H(X_{[n]} | Y_{[n]}) &= H(X_{[2,n-1]} | X_1, X_n, Y_{[n]}) + H(X_1, X_n | Y_{[n]}) \\ &\leq 0 + H(X_1, X_n | Y_1, Y_n) \\ &\leq 2H(X_1 | Y_1) \\ &\leq 2 \max \{ \text{ld } |g^{-1}(y)| : y \in \mathcal{Y} \} \end{aligned}$$

$$\leq 2 \operatorname{ld}(|\mathcal{X}| - |\mathcal{Y}| + 1).$$

□

*Proof of (11b).* As  $\mathcal{K} = \infty$ , the bound from (6) holds uniformly. Thus

$$\overline{H}(X|Y) = \lim_{n \rightarrow \infty} \frac{1}{n} H(X_{[n]}|Y_{[n]}) \leq \lim_{n \rightarrow \infty} \frac{2 \operatorname{ld}(|\mathcal{X}| - |\mathcal{Y}| + 1)}{n} = 0.$$

□

*Proof of (12b).* Recall that we assume  $\mathcal{K} = \infty$ . We show that, for  $\mathbf{y} \in \mathcal{Y}^n$  with  $\mathbb{P}(Y_{[n]} = \mathbf{y}) > 0$ , we have

$$\mathbb{P}(T_n \leq (|\mathcal{X}| - |\mathcal{Y}| + 1)^2 | Y_{[n]} = \mathbf{y}) = 1. \quad (15)$$

This implies (12b). To show (15), we use (13) to bound

$$\begin{aligned} & \sum_{\mathbf{x} \in g^{-1}(\mathbf{y})} [\mathbb{P}(X_{[n]} = \mathbf{x}) > 0] \\ &= \sum_{x_1, x_n \in g^{-1}(\mathbf{y}_{\{1, n\}})} [\mathbb{P}(X_1 = x_1, X_n = x_n | Y_{[n]} = \mathbf{y}) > 0] \\ & \quad \times \sum_{\mathbf{x} \in g^{-1}(\mathbf{y}_{[2, n-1]})} [\mathbb{P}(X_{[2, n-1]} = \mathbf{x} | X_1 = x_1, X_n = x_n, Y_{[2, n-1]} = \mathbf{y}_{[2, n-1]}) > 0] \\ &\leq \sum_{x_1, x_n \in g^{-1}(\mathbf{y}_{\{1, n\}})} [\mathbb{P}(X_1 = x_1, X_n = x_n | Y_{[n]} = \mathbf{y}) > 0] \\ &\leq |g^{-1}(\mathbf{y}_{\{1, n\}})| \leq (|\mathcal{X}| - |\mathcal{Y}| + 1)^2. \end{aligned}$$

□

### 3.2 Non-overlapping traversal instants

The main result of this section in proposition 11 is an almost-sure *linear lower growth bound* for non-overlapping occurrences of a fixed, finite pattern in a realisation.

Let  $Z := (Z_n)_{n \in \mathbb{N}}$  be a stationary stochastic process taking values in  $\mathcal{Z}$ . The *occupation instants* of a state  $z$  is the set of indices

$$\mathcal{O}_Z^z(n) := \{i \in [n] : Z_i = z\}. \quad (16a)$$

The classic *occupation time* [Par99, section 6.4] is the cardinality of the occupation instants. The *traversal instants* of a sequence  $\mathbf{z} \in \mathcal{Z}^k$  is the set of indices

$$\mathcal{T}_Z^{\mathbf{z}}(n) := \{i \in [n - k + 1] : Z_{[i, i+k-1]} = \mathbf{z}\}. \quad (16b)$$

The *non-overlapping traversal instants* of a sequence  $\mathbf{z} \in \mathcal{Z}^k$  is the set of indices

$$\mathcal{N}_Z^{\mathbf{z}}(n) := \left\{ i \in [n - k + 1] : \begin{array}{l} Z_{[i, i+k-1]} = \mathbf{z} \\ \forall j \in [i + 1, i + k - 1] : Z_{[j, j+k-1]} \neq \mathbf{z} \end{array} \right\}, \quad (16c)$$

where we select lower indices greedily.

For  $k \in \mathbb{N}$ , the  $k$ -transition process  $Z^{(k)}$  of  $Z$  is the stochastic process on  $\mathcal{Z}^k$  with marginals

$$\mathbb{P}(Z_{[n]}^{(k)} = (\mathbf{z}^i)_{i=1}^n) = \mathbb{P}(Z_{[n-1]} = (\mathbf{z}_{\{1\}}^i)_{i=1}^{n-1}, Z_{[n, n+k-1]} = \mathbf{z}^n), \quad (17)$$

if  $\forall i \in [n-1] : \mathbf{z}_{[2, k]}^i = \mathbf{z}_{[k-1]}^{i+1}$ , and zero else. Obvious relations are

$$\mathcal{T}_Z^{\mathbf{z}}(n) = \mathcal{O}_{Z^{(k)}}^{\mathbf{z}}(n-k) \quad (18a)$$

and

$$\mathcal{N}_Z^{\mathbf{z}}(n) \subseteq \mathcal{T}_Z^{\mathbf{z}}(n) \quad \text{with} \quad |\mathcal{N}_Z^{\mathbf{z}}(n)| \geq \frac{1}{k} |\mathcal{T}_Z^{\mathbf{z}}(n)|. \quad (18b)$$

**Proposition 11.** Let  $\mathbf{s} \in \mathcal{X}^k$  with  $p := \mathbb{P}(X_{[k]} = \mathbf{s} | X_1 = \mathbf{s}_{\{1\}}) > 0$ . Then

$$\mathbb{P} \left( \liminf_{n \rightarrow \infty} \frac{1}{n} |\mathcal{N}_X^{\mathbf{s}}(n)| \geq \frac{p\mu(\mathbf{s}_{\{1\}})}{k} \right) = 1 \quad (19a)$$

and

$$\forall \varepsilon > 0 : \quad \lim_{n \rightarrow \infty} \mathbb{P} \left( |\mathcal{N}_X^{\mathbf{s}}(n)| \geq \left( \frac{p\mu(\mathbf{s}_{\{1\}})}{k} - \varepsilon \right) n \right) = 1. \quad (19b)$$

**Lemma 12** (Ergodic theorem [Woe09, theorem 3.55 on page 69]). *For every homogeneous, irreducible and aperiodic Markov chain  $Z := (Z_n)_{n \in \mathbb{Z}}$  on a finite state space  $\mathcal{Z}$  with invariant measure  $\nu$ , all  $f : \mathcal{Z} \rightarrow \mathbb{R}$  and each starting distribution  $\alpha \in \mathcal{M}_1(\mathcal{Z})$  of  $Z_1$ , we have*

$$\mathbb{P}_\alpha \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n f(Z_i) = \int_{\mathcal{Z}} f(z) d\nu(z) =: \nu(f) \right) = 1. \quad (20)$$

*Proof of proposition 11.* Statement (19b) is a direct consequence of (19a).

The  $k$ -transition process  $X^{(k)}$  is a Markov chain with transition probabilities

$$P_{(\mathbf{x}, \mathbf{x}')}^{(k)} := \begin{cases} P_{(\mathbf{x}_{\{k\}}, \mathbf{x}'_{\{k\}})} & \text{if } \mathbf{x}_{[2, k]} = \mathbf{x}'_{[k-1]}, \\ 0 & \text{else.} \end{cases} \quad (21)$$

Furthermore, as  $X$  is irreducible and aperiodic, then so is  $X^{(k)}$ . Its invariant measure  $\mu^{(k)}$  fulfils  $\mu^{(k)}(\mathbf{x}) = \mu(\mathbf{x}_{\{1\}}) \prod_{i=1}^{k-1} P_{(\mathbf{x}_{\{i\}}, \mathbf{x}_{\{i+1\}})}$ .

Let  $f$  be the indicator function of  $\mathbf{s}$ . We use (18) and lemma 12 to derive

$$\begin{aligned} & \mathbb{P} \left( \liminf_{n \rightarrow \infty} \frac{1}{n} |\mathcal{N}_X^{\mathbf{s}}(n)| \geq \frac{\mu^{(k)}(f)}{k} \right) \\ & \geq \mathbb{P} \left( \liminf_{n \rightarrow \infty} \frac{1}{n} |\mathcal{T}_X^{\mathbf{s}}(n)| \geq \mu^{(k)}(f) \right) \\ & = \mathbb{P} \left( \liminf_{n \rightarrow \infty} \frac{1}{n} |\mathcal{O}_{X^{(k)}}^{\mathbf{s}}(n-k)| \geq \mu^{(k)}(f) \right) \\ & = \mathbb{P} \left( \lim_{n \rightarrow \infty} \frac{1}{n} |\mathcal{O}_{X^{(k)}}^{\mathbf{s}}(n)| \geq \mu^{(k)}(f) \right) \\ & = 1. \end{aligned}$$

Finally  $\mu^{(k)}(f) = \mu^{(k)}(\mathbf{s}) = p\mu(\mathbf{s}_{\{1\}})$ . □

### 3.3 Conditional Markov property

This section presents two technical statements about discrete Markov processes. Let  $X := (X_n)_{n \in \mathbb{Z}}$  be a stochastic process on the Cartesian product  $\mathcal{S} := \prod_{n \in \mathbb{Z}} \mathcal{S}_n$  of the finite sets  $(\mathcal{S}_n)_{n \in \mathbb{Z}}$ . For  $A \subseteq \mathbb{Z}$  let  $\mathcal{S}_A := \prod_{n \in A} \mathcal{S}_n$ . In the remainder of this section, we assume that all conditional probabilities are well-defined. The process  $X$  is Markov, iff

$$\begin{aligned} \forall n \in \mathbb{Z}, m \in \mathbb{N}, s_n \in \mathcal{S}_n, s_{[n-m, n-1]} \in \mathcal{S}_{[n-m, n-1]} : \\ \mathbb{P}(X_n = s_n | X_{[n-m, n-1]} = s_{[n-m, n-1]}) = \mathbb{P}(X_n = s_n | X_{n-1} = s_{n-1}). \end{aligned} \quad (22)$$

We denote by  $A \Subset \mathbb{Z}$  the fact that  $A$  is a finite subset of  $\mathbb{Z}$ . The first statement is a factorisation of conditional probabilities over disjoint index blocks:

$$\begin{aligned} \forall m \in \mathbb{N}, \emptyset \neq A_1, B_1, \dots, B_{m-1}, A_m \Subset \mathbb{Z}, B_0, B_m \Subset \mathbb{Z}, \\ A \cap B = \emptyset \text{ where } A := \biguplus_{i=1}^m A_i \text{ and } B := \biguplus_{i=0}^m B_i, x_A \in \mathcal{S}_A, x_B \in \mathcal{S}_B, \\ (\forall i \in [m] : b_i^- := \sup(B_{i-1}), b_i^+ := \inf(B_i), A_i \subseteq ]b_i^-, b_i^+[) : \\ \mathbb{P}(X_A = x_A | X_B = x_B) = \prod_{i=1}^m \mathbb{P}(X_{A_i} = x_{A_i} | X_{b_i^-} = x_{b_i^-}, X_{b_i^+} = x_{b_i^+}). \end{aligned} \quad (23)$$

Secondly, a Markov process retains the Markov property under a Cartesian conditioning:

$$\forall \emptyset \neq C \Subset \mathbb{Z}, S_C := \prod_{n \in C} \mathcal{S}_n \text{ with } S_n \subseteq \mathcal{S}_n : (X | X_C \in S_C) \text{ is Markov.} \quad (24)$$

*Proof.* We need the intermediate statements

$$\begin{aligned} \forall n \in \mathbb{Z}, \emptyset \neq B \Subset \mathbb{Z} : -\infty, n[ \cap B = \emptyset, n \in B : \\ \mathbb{P}(X_n = x_n | X_B = x_B) = \mathbb{P}(X_n = x_n | X_{\max(B)} = x_{\max(B)}) \end{aligned} \quad (25)$$

and

$$\begin{aligned} \forall \emptyset \neq A, B \Subset \mathbb{Z}, \max(B) < \min(A), x_A \in \mathcal{S}_A, S_B \subseteq \{x_{\max(B)}\} \times \mathcal{S}_{B \setminus \max(B)} : \\ \mathbb{P}(X_A = x_A | X_B \in S_B) = \mathbb{P}(X_A = x_A | X_{\max(B)} = x_{\max(B)}). \end{aligned} \quad (26)$$

Proof of (25): Let  $C := ]\max(B), n[$  and  $D := [\min(B), \max(B)] \setminus B$ . We use (22) to get

$$\begin{aligned} & \mathbb{P}(X_n = x_n | X_B = x_B) \\ &= \frac{\sum_{x_C, x_D} \mathbb{P}(X_n = x_n, X_C = x_C, X_B = x_B, X_D = x_D)}{\mathbb{P}(X_B = x_B)} \\ &= \frac{\sum_{x_C, x_D} \mathbb{P}(X_n = x_n, X_C = x_C | X_B = x_B, X_D = x_D) \mathbb{P}(X_B = x_B, X_D = x_D)}{\mathbb{P}(X_B = x_B)} \\ &= \frac{\sum_{x_C} \mathbb{P}(X_n = x_n, X_C = x_C | X_{\max(B)} = x_{\max(B)}) \sum_{x_D} \mathbb{P}(X_B = x_B, X_D = x_D)}{\mathbb{P}(X_B = x_B)} \end{aligned}$$

$$= \mathbb{P}(X_n = x_n | X_{\max(B)} = x_{\max(B)}).$$

Proof of (23): For  $A \in \mathbb{N}$ , we abbreviate the event  $E_A := (X_A = x_A)$ . Apply (25) to get

$$\begin{aligned} & \mathbb{P}(X_A = x_A | X_B = x_B) \\ &= \frac{\mathbb{P}(X_A = x_A, X_B = x_B)}{\mathbb{P}(X_B = x_B)} \\ &= \frac{\mathbb{P}(E_{B_0}) \prod_{i=1}^m \mathbb{P}(E_{B_i \setminus \{b_i^+\}} | (E_{A_j})_{j \leq i}, (E_{B_j})_{j < i}, E_{b_i^+}) \mathbb{P}(E_{b_i^+}, E_{A_i} | (E_{A_j})_{j < i}, (E_{B_j})_{j < i})}{\mathbb{P}(E_{B_0}) \prod_{i=1}^m \mathbb{P}(E_{B_i \setminus \{b_i^+\}} | (E_{B_j})_{j < i}, E_{b_i^+}) \mathbb{P}(E_{b_i^+} | (E_{B_j})_{j < i})} \\ &= \prod_{i=1}^m \frac{\mathbb{P}(E_{B_i \setminus \{b_i^+\}} | E_{b_i^+}) \mathbb{P}(E_{b_i^+}, E_{A_i} | E_{b_i^-})}{\mathbb{P}(E_{B_i \setminus \{b_i^+\}} | E_{b_i^+}) \mathbb{P}(E_{b_i^+} | E_{b_i^-})} \\ &= \prod_{i=1}^m \mathbb{P}(E_{A_i} | E_{b_i^-}, E_{b_i^+}). \end{aligned}$$

Proof of (26): Let  $b := \max(B)$ . We apply (25) to get

$$\begin{aligned} & \mathbb{P}(X_A = x_A | X_B \in S_B) \\ &= \frac{\sum_{x_B \in S_B} \mathbb{P}(X_A = x_A, X_B = x_B)}{\mathbb{P}(X_B \in S_B)} \\ &= \frac{\sum_{x_B \in S_B} \mathbb{P}(X_A = x_A | X_b = x_b) \mathbb{P}(X_B = x_B)}{\mathbb{P}(X_B \in S_B)} \\ &= \mathbb{P}(X_A = x_A | X_b = x_b). \end{aligned}$$

Proof of (24): Let  $n \in \mathbb{Z}$ ,  $m \in \mathbb{N}$ ,  $B := [n - m, n[$ ,  $x_n \in \mathcal{S}_n$ ,  $x_B \in \mathcal{S}_B$ . Let  $C_+ := C \cap [n, \infty[$  and  $C_- := C \cap ]-\infty, n[$ . Thus  $S_C = S_{C_-} \times S_{C_+}$ . We apply (26) twice to show that  $(X | X_C \in S_C)$  fulfils (22) and is thus Markov:

$$\begin{aligned} & \mathbb{P}(X_n = x_n | X_B = x_B, X_C \in S_C) \\ &= \frac{\mathbb{P}(X_n = x_n, X_{C_+} \in S_{C_+}, X_B = x_B, X_{C_-} \in S_{C_-})}{\mathbb{P}(X_B = x_B, X_C \in S_C)} \\ &= \frac{\mathbb{P}(X_n = x_n, X_{C_+} \in S_{C_+} | X_B = x_B, X_{C_-} \in S_{C_-}) \mathbb{P}(X_B = x_B, X_{C_-} \in S_{C_-})}{\mathbb{P}(X_{C_+} \in S_{C_+} | X_B = x_B, X_{C_-} \in S_{C_-}) \mathbb{P}(X_B = x_B, X_{C_-} \in S_{C_-})} \\ &= \frac{\mathbb{P}(X_n = x_n, X_{C_+} \in S_{C_+} | X_{n-1} = x_{n-1})}{\mathbb{P}(X_{C_+} \in S_{C_+} | X_{n-1} = x_{n-1})} \\ &= \mathbb{P}(X_n = x_n | X_{n-1} = x_{n-1}, X_{C_+} \in S_{C_+}) \\ &= \mathbb{P}(X_n = x_n | X_{n-1} = x_{n-1}, X_{C_+} \in S_{C_+}, X_{C_-} \in S_{C_-}) \\ &= \mathbb{P}(X_n = x_n | X_{n-1} = x_{n-1}, X_C \in S_C). \end{aligned}$$

□

### 3.4 The loss case

We start with some derivations common to the proof of (11a) and (12a). We assume  $\mathcal{K} < \infty$ . Equation (3) is equivalent to the existence of  $\tilde{x}, \hat{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}^{\mathcal{K}}, \mathbf{x} \in g^{-1}(\mathbf{y})$  with

$$0 < \mathbb{P}(X_0 = \tilde{x}, X_{[\mathcal{K}]} = \mathbf{x}, X_{\mathcal{K}+1} = \hat{x}) < \mathbb{P}(X_0 = \tilde{x}, Y_{[\mathcal{K}]} = \mathbf{y}, X_{\mathcal{K}+1} = \hat{x}). \quad (27)$$

Let  $\mathbf{s} := (\tilde{x}, \mathbf{x}, \hat{x})$ . The *unreconstructable set of trajectories*  $\mathcal{H}$  is

$$\mathcal{H} := \{\tilde{x}\} \times g^{-1}(\mathbf{y}) \times \{\hat{x}\}. \quad (28)$$

Equation (3) implies that  $\mathcal{H}$  contains at least two elements with positive probability. If we pass through  $\mathcal{H}$ , we incur an *entropy loss*  $L$ :

$$L := H(X_{[\mathcal{K}]} | X_{[0, \mathcal{K}+1]} \in \mathcal{H}) > 0. \quad (29)$$

Let  $\mathcal{I}$  be the random set of indices marking the start of non-overlapping runs of  $X_{[n]}$  through  $\mathcal{H}$ , that is

$$\mathcal{I} := \left\{ i \in [n - \mathcal{K} - 1] : \begin{array}{c} X_{[i, i+\mathcal{K}+1]} \in \mathcal{H} \\ \text{and} \\ \forall j \in [i+1, i+\mathcal{K}+1] : X_{[j, j+\mathcal{K}+1]} \notin \mathcal{H} \end{array} \right\}, \quad (30)$$

where we select lower indices greedily. For the  $\mathbf{s}$  from after (27), we lower-bound the tail probability of the cardinality of  $\mathcal{I}$  by the one of  $\mathcal{N}_X^{\mathbf{s}}(n)$ :

$$\forall m \in \mathbb{N} : \quad \mathbb{P}(|\mathcal{I}| \geq m) \geq \mathbb{P}(|\mathcal{N}_X^{\mathbf{s}}(n)| \geq m). \quad (31)$$

Finally, let

$$\alpha := \frac{\mathbb{P}(X_{[\mathcal{K}+2]} = \mathbf{s})}{2(\mathcal{K} + 2)} > 0. \quad (32)$$

*Proof of (11a).* We claim that, for every  $m \in \mathbb{N}$ :

$$H(X_{[n]} | Y_{[n]}) \geq \mathbb{P}(|\mathcal{I}| \geq m) H(X_{[n]} | Y_{[n]}, |\mathcal{I}| \geq m) \geq \mathbb{P}(|\mathcal{I}| \geq m) mL. \quad (33)$$

Combining (33) and (31), for  $m = \alpha n$ , with (19b), we arrive at (11a):

$$\begin{aligned} \overline{H}(X|Y) &= \lim_{n \rightarrow \infty} \frac{1}{n} H(X_{[n]} | Y_{[n]}) \\ &\geq \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{P}(|\mathcal{I}| \geq \alpha n) \alpha n L \geq \alpha L \lim_{n \rightarrow \infty} \mathbb{P}(|\mathcal{N}_X^{\mathbf{s}}(n)| \geq \alpha n) = \alpha L > 0. \end{aligned} \quad (34)$$

It rests to prove (33). We fix  $m, n \in \mathbb{N}$ . For  $I \subseteq [n]$  with  $\mathbb{P}(\mathcal{I} = I) > 0$  and each  $i \in I$ , we derive the indices of the block  $B_i := [i, i + \mathcal{K} + 1]$  and its interior  $\widehat{B}_i := [i + 1, i + \mathcal{K}]$ . Their unions are  $B := \biguplus_{i \in I} B_i$  and  $\widehat{B} := \biguplus_{i \in I} \widehat{B}_i$  respectively. Hence

$$\begin{aligned} &H(X_{[n]} | Y_{[n]}, \mathcal{I} = I), \\ &\geq H(X_{\widehat{B}} | X_{[n] \setminus B}, \forall i \in I : X_{B_i} \in \mathcal{H}), \end{aligned} \quad (35a)$$

$$= H(X_{\widehat{B}} | \forall i \in I : X_{B_i} \in \mathcal{H}), \quad (35b)$$

$$= \sum_{i \in I} H(X_{\widehat{B}_i} | X_{B_i} \in \mathcal{H}), \quad (35c)$$

$$= |I| \times L, \quad (35d)$$

where in (35a) we throw away all information outside  $\widehat{B}$  and condition on it, in (35b) we apply the conditional factorisation (23) twice to remove every condition except the block ends, in (35c) we apply the conditional factorisation (23) to the Markov process  $(X | X_B \in \mathcal{H}^{|I|})$  and in (35d) we conclude by stationarity and the minimum loss (29). Hence

$$\begin{aligned} H(X_{[n]} | Y_{[n]}, |\mathcal{I}| \geq m) &= \sum_{\substack{I \subseteq [n] \\ |I| \geq m}} \mathbb{P}(\mathcal{I} = I | |\mathcal{I}| \geq m) H(X_{[n]} | Y_{[n]}, \mathcal{I} = I) \\ &\geq \sum_{\substack{I \subseteq [n] \\ |I| \geq m}} \mathbb{P}(\mathcal{I} = I | |\mathcal{I}| \geq m) \times |I| \times L \\ &\geq mL. \end{aligned}$$

□

*Proof of (12a).* For the  $\mathbf{s}$  from after (27), we  $\mathbb{P}$ -almost surely have

$$T_n \geq 2^{|\mathcal{N}_X^{\mathbf{s}}(n)|}. \quad (36)$$

Thus, (36) and (19a) imply that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \sqrt[n]{T_n} &\geq \liminf_{n \rightarrow \infty} \exp((\log 2) \frac{1}{n} |\mathcal{N}_X^{\mathbf{s}}(n)|) \\ &= \exp((\log 2) \liminf_{n \rightarrow \infty} \frac{1}{n} |\mathcal{N}_X^{\mathbf{s}}(n)|) \geq \exp((\log 2)\alpha) = 2^\alpha > 1. \end{aligned} \quad (37)$$

□

*Proof of proposition 3.* Let  $x_0, x_{\mathcal{K}+1}, \mathbf{y}, \mathbf{x}', \mathbf{x}''$  as in (3). Suppose that  $\mathcal{K} > K := \sum_{y \in \mathcal{Y}} |g^{-1}(y)|(|g^{-1}(y)| - 1)$  and  $\mathcal{K} > 1$ . We apply the *pigeon-hole principle* first to every  $x \in g^{-1}(y)$  and then to each  $g^{-1}(y)$ , for every  $y \in \text{supp } \mathbf{y}$ . This ensures that the two trajectories intersect:

$$\exists m \in [\mathcal{K}] : \quad \mathbf{x}'_{\{m\}} = \mathbf{x}''_{\{m\}}. \quad (38)$$

Choose  $m$  fulfilling (38). If  $m = 1$ , then  $\mathbf{x}'_{\{1\}}, x_{\mathcal{K}+1}, \mathbf{y}_{[2, \mathcal{K}]}, \mathbf{x}'_{[2, \mathcal{K}]}, \mathbf{x}''_{[2, \mathcal{K}]}$  fulfil the conditions in (3). If  $m > 1$ , then  $x_0, \mathbf{x}'_{\{m\}}, \mathbf{y}_{[m-1]}, \mathbf{x}'_{[m-1]}, \mathbf{x}''_{[m-1]}$  fulfil the conditions in (3). Both cases lead to  $\mathcal{K} < \mathcal{K}$ , a contradiction. □

## 4 Proofs of the sufficient conditions

### 4.1 Single entry implies entropy rate preservation

*Proof of proposition 6.*

$$\overline{H}(X|Y)$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \frac{1}{n} H(X_{[n]} | Y_{[n]}) \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n H(X_i | X_{[i-1]}, Y_{[n]}) \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n H(X_i | X_{[i-1]}, Y_i) \tag{39a}
\end{aligned}$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n H(X_i | X_{i-1}, Y_i) \tag{39b} \\
&= H(X_1 | X_0, Y_1)
\end{aligned}$$

where (39a) is due to  $Y_{[i-1]} = g(X_{[i-1]})$  and because conditioning reduces entropy and (39b) is by Markovity of  $X$  from (24). We continue

$$H(X_1 | X_0, Y_1) = - \sum_{\substack{y \in \mathcal{Y} \\ x_1 \in g^{-1}(y) \\ x_0 \in \mathcal{X}}} \mathbb{P}(X_1 = x_1, X_0 = x_0) \text{ld} \frac{\mathbb{P}(X_1 = x_1 | X_0 = x_0)}{\mathbb{P}(Y_1 = y | X_0 = x_0)} = 0,$$

because  $(P, g)$  is SE. Thus  $\mathbb{P}(X_1 = x_1 | X_0 = x_0) = \mathbb{P}(Y_1 = y | X_0 = x_0)$ , if  $x_1 = x'(x_0, y)$ , and zero otherwise.  $\square$

## 4.2 Single forward property implies $k$ -lumpability

For (conditional) probabilities we use the following short-hand notation:

$$\mathbb{P}(Z = z) = p_Z(z) \quad \text{and} \quad \mathbb{P}(Z_1 = z_1 | Z_2 = z_2) = p_{Z_1 | Z_2}(z_1 | z_2),$$

where we always assume that the latter is well-defined, i.e., that  $p_{Z_2}(z_2) > 0$ .

We start with some auxiliary results:

**Proposition 13.**

$$Y \text{ is HMC}(k) \Leftrightarrow \overline{H}(Y) = H(Y_k | Y_{[0, k[}). \tag{40}$$

*Proof.*

$$\begin{aligned}
0 &= H(Y_k | Y_{[0, k[}) - \overline{H}(Y) \\
&= \lim_{n \rightarrow \infty} H(Y_k | Y_{[0, k[}) - H(Y_n | Y_{[0, n[}) \\
&= \lim_{n \rightarrow \infty} H(Y_n | Y_{[n-k, n[}) - H(Y_n | Y_{[0, n[}) \\
&= \lim_{n \rightarrow \infty} I(Y_n; Y_{[0, n-k[} | Y_{[n-k, n[}),
\end{aligned}$$

where the last quantity is the *conditional mutual information* between  $Y_n$  and  $Y_{[0, n-k[}$  conditioned on  $Y_{[n-k, n[}$ . By stationarity, this sequence increases monotonically in  $n$ , thus, setting it to zero implies [Gra90, Lemma 3.15, pp. 88] that for all  $n \in \mathbb{N}$ :

$$\begin{aligned}
&p_{Y_n | Y_{[n-k, n[}}(\cdot | \mathbf{y}) p_{Y_{[0, n-k[} | Y_{[n-k, n[}}(\cdot | \mathbf{y}) \\
&= p_{Y_n, Y_{[0, n-k[} | Y_{[n-k, n[}}(\cdot | \mathbf{y})
\end{aligned}$$

$$= p_{Y_n|Y_{[0,n[}}(\cdot|\cdot, \mathbf{y}) p_{Y_{[0,n-k[}|Y_{[n-k,n[}}(\cdot|\mathbf{y}),$$

where the first equality holds  $p_{Y_{[n-k,n[}}$ -a.s. The equality between the first and last line is equivalent to the Markov property in definition 8.  $\square$

**Lemma 14** ([CT06, Thm. 4.5.1, pp. 86]). *If  $X$  is a stationary Markov chain and  $Y$  a stationary process defined by  $Y_n := g(X_n)$ , then the entropy rate of  $Y$  is bounded by*

$$\forall k \in \mathbb{N}: \quad H(Y_k|Y_{[k-1]}, X_0) \leq \overline{H}(Y) \leq H(Y_k|Y_{[0,k[}].$$

A direct solution for the entropy rate of the lumped process  $Y$  was shown to be intrinsically complicated in [Bla57]. However, both the upper and lower bound are asymptotically tight [CT06, Thm. 4.5.1, pp. 86].

**Proposition 15.** *The following facts are equivalent:*

$$\forall \mathbf{y} \in \mathcal{Y}^k, y \in \mathcal{Y}, x \in g^{-1}(y):$$

$$p_{Y_k|Y_{[1,k[}, X_0}(y|\mathbf{y}, x) = p_{Y_k|Y_{[1,k[}, Y_0}(y|\mathbf{y}, g(x)), \quad (41a)$$

$$H(Y_k|Y_{[k-1]}, X_0) = H(Y_k|Y_{[0,k[}). \quad (41b)$$

*They are sufficient for  $k$ -lumpability of  $(P, g)$ , but necessary only for strongly  $k$ -lumpable  $(P, g)$ .*

*Proof.* Statement (41b), together with proposition 13 and lemma 14, implies the  $k$ -lumpability of  $(P, g)$ .

Equivalence between (41b) and (41a):

$$\begin{aligned} 0 &= H(Y_k|Y_{[0,k[}) - H(Y_k|Y_{[k-1]}, X_0) \\ &= H(Y_k|Y_{[0,k[}) - H(Y_k|Y_{[0,k[}, X_0) \\ &= I(Y_k; X_0|Y_{[0,k[}), \end{aligned}$$

which is equivalent to

$$p_{Y_k, X_0|Y_{[0,k[}}(\cdot|\mathbf{y}) = p_{Y_k|Y_{[0,k[}}(\cdot|\mathbf{y}) p_{X_0|Y_{[0,k[}}(\cdot|\mathbf{y})$$

and thus

$$p_{Y_k|Y_{[0,k[}, X_0}(\cdot|\mathbf{y}, x) = p_{Y_k|Y_{[0,k[}}(\cdot|\mathbf{y}),$$

for all  $x, \mathbf{y}$  such that  $p_{X_0, Y_{[0,k[}}(x, \mathbf{y}) > 0$ .

The proof that strong  $k$ -lumpability implies (41) follows the one in [KS76, Thm. 6.3.2]: Strong lumpability requires that  $p_{Y_k|Y_{[0,k[}}(y|\mathbf{y})$  is independent of the distribution of  $X_0$ , thus in particular, it needs to have the same value for all distributions placing unit mass on a state in  $x' \in g^{-1}(\mathbf{y}_{\{0\}})$ . We have

$$\begin{aligned} &p_{Y_k|Y_{[0,k[}}(y|\mathbf{y}) \\ &= \sum_{x \in g^{-1}(\mathbf{y}_{\{0\}})} p_{Y_k|Y_{[0,k[}, X_0}(y|\mathbf{y}, x) p_{X_0|Y_{[0,k[}}(x|\mathbf{y}) \end{aligned} \quad (42)$$

Since

$$p_{X_0|Y_{[0,k[}}(x|\mathbf{y}) = \frac{p_{X_0}(x)p_{Y_{[0,k[}|X_0}(\mathbf{y}|x)}{p_{Y_{[0,k[}}(\mathbf{y})}$$

and since

$$\sum_{x \in g^{-1}(\mathbf{y}_{\{0\}})} p_{X_0|Y_{[0,k[}}(x|\mathbf{y}) = 1,$$

it follows that  $p_{Y_{[0,k[}|X_0}(\mathbf{y}|x') = p_{Y_{[0,k[}}(\mathbf{y})$  ( $p_{X_0}(x) = 0$ , for all  $x \in g^{-1}(\mathbf{y}_{\{0\}}) \setminus \{x'\}$ ). The sum in (42) degenerates and we obtain

$$p_{Y_k|Y_{[0,k[}}(y|\mathbf{y}) = p_{Y_k|Y_{[0,k[,X_0}}(y|\mathbf{y}, x').$$

Strong lumpability implies that  $p_{Y_k|Y_{[0,k[}}(y|\mathbf{y})$  is independent of the distribution of  $X_0$ , whence (41) follows.  $\square$

We are currently not sure if (41) and strong  $k$ -lumpability are equivalent.

**Example 16** (taken from [KS76, pp. 139]). *Consider the following transition matrix, where the lines divide lumped states:*

$$P := \left[ \begin{array}{c|ccc} 1/4 & 1/16 & 3/16 & 1/2 \\ \hline 0 & 1/12 & 1/12 & 5/6 \\ 0 & 1/12 & 1/12 & 5/6 \\ 7/8 & 1/32 & 3/32 & 0 \end{array} \right].$$

*This lumping (and its time-reversal) is 1-lumpable [KS76, pp. 139]. However, we have (with an accuracy of 0.0001)*

$$0.5588 = H(Y_1|X_0) < \overline{H}(Y) = H(Y_1|Y_0) = 0.9061$$

and

$$0.9048 = H(Y_0|X_1) < \overline{H}(\hat{Y}) = H(Y_0|Y_1) = 0.9061.$$

Hence  $k$ -lump does not imply (41).

*Proof of proposition 10.* We first show that SE contains SFS( $k$ ), which implies preservation of entropy. We have

$$p_{X_{[k-1]}|Y_{[k-1]},X_0}(\mathbf{x}|\mathbf{y}_{[k-1]},x_0) = \begin{cases} 1 & \text{if } \mathbf{x} = \mathbf{x}'(y_{[0,k-1]}) \text{ from (10),} \\ 0 & \text{else.} \end{cases}$$

If SE does not hold, then there exist states  $y^* \in \mathcal{Y}$  and  $x^* \in \mathcal{X}$  such that at least two states in  $g^{-1}(y^*)$  have positive transition probabilities from  $x^*$ . Assume that  $\mathbf{y}_{\{k-2,k-1\}} = (g(x^*), y^*)$ . Thus, observing  $\mathbf{y}_{[0,k[}$  does not determine  $\mathbf{x}$ .

Second, we show that SFS( $k$ ) implies  $k$ -lumpability of  $(P, g)$ . By proposition 15, it suffices to show that

$$p_{Y_k|Y_{[k-1]},X_0}(\mathbf{y}_{\{k\}}|\mathbf{y}_{[k-1]},x_0) = p_{Y_k|Y_{[0,k[}}(\mathbf{y}_{\{k\}}|\mathbf{y}_{[0,k[})$$

holds  $\mathbb{P}$ -almost surely. Knowing  $Y_{[0,k[} = \mathbf{y}_{[0,k[}$ , however, determines  $X_{[k-1]}$  regardless of  $x_0 \in g^{-1}(y_0)$ . The Markov property implies

$$\begin{aligned}
p_{Y_k|Y_{[k-1]}, X_0}(\mathbf{y}_{\{k\}}|\mathbf{y}_{[k-1]}, x_0) &= p_{Y_k|Y_{[0,k[}}(\mathbf{y}_{\{k\}}|\mathbf{y}_{[0,k[}) \\
&= p_{Y_k|X_{k-1}}(\mathbf{y}_{\{k\}}|\mathbf{x}'(\mathbf{y}_{[0,k[})_{\{k-1\}}).
\end{aligned}$$

Moreover, the SE property implies that, for every  $x \in \mathcal{X}, y \in \mathcal{Y}$ ,

$$p_{Y_k|X_{k-1}}(y|x) = p_{X_k|X_{k-1}}(x'(x, y)|x),$$

where  $x'(x, y)$  is from (8). This yields another proof for entropy rate preservation by equating the outer terms of the following chain of inequalities (the first inequality is due to lemma 14, the second due to data processing [GK11, WA60]):

$$H(Y_k|Y_{[k-1]}, X_0) \leq \overline{H}(Y) \leq \overline{H}(X) = H(X_k|X_{k-1}).$$

□

## References

- [And99] Brian D. O. Anderson. The realization problem for hidden Markov models. *Math. Control Signals Systems*, 12(1):80–120, 1999.
- [BdM<sup>+</sup>92] Peter F. Brown, Peter V. deSouza, Robert L. Mercer, Vincent J. Della Pietra, and Jenifer C. Lai. Class-based n-gram models of natural language. *Comput. Linguist.*, 18(4):467–479, December 1992.
- [BK57] David Blackwell and Lambert Koopmans. On the identifiability problem for functions of finite Markov chains. *Ann. Math. Statist.*, 28:1011–1015, 1957.
- [Bla57] David Blackwell. The entropy of functions of finite-state Markov chains. In *Transactions of the first Prague conference on information theory, Statistical decision functions, random processes held at Liblice near Prague from November 28 to 30, 1956*, pages 13–20. Publishing House of the Czechoslovak Academy of Sciences, Prague, 1957.
- [BR58] C. J. Burke and M. Rosenblatt. A Markovian function of a Markov chain. *Ann. Math. Statist.*, 29:1112–1122, 1958.
- [Car67] J. W. Carlyle. Identification of state-calculable functions of finite Markov chains. *Ann. Math. Statist.*, 38:201–205, 1967.
- [CT06] Thomas M. Cover and Joy A. Thomas. *Elements of information theory*. Wiley-Interscience [John Wiley & Sons], Hoboken, NJ, second edition, 2006.
- [EM02] Yariv Ephraim and Neri Merhav. Hidden Markov processes. *IEEE Trans. Inform. Theory*, 48(6):1518–1569, 2002. Special issue on Shannon theory: perspective, trends, and applications.
- [Gil59] Edgar J. Gilbert. On the identifiability problem for functions of finite Markov chains. *Ann. Math. Statist.*, 30:688–697, 1959.
- [GK11] Bernhard Claus Geiger and Gernot Kubin. Some results on the information loss in dynamical systems. In *Proc. IEEE Int. Sym. Wireless Communication Systems (ISWSC)*, pages 794–798, Aachen, November 2011. extended version available: [arXiv:1106.2404](https://arxiv.org/abs/1106.2404) [cs.IT].
- [GL05] Leonid Gurvits and James Ledoux. Markov property for a function of a Markov chain: a linear algebra approach. *Linear Algebra Appl.*, 404:85–117, 2005.
- [Gra90] Robert M. Gray. *Entropy and information theory*. Springer-Verlag, New York, 1990.
- [HMMW10] Thomas A. Henzinger, Linar Mikeev, Maria Mateescu, and Verena Wolf. Hybrid numerical solution of the chemical master equation.

- In *Proceedings of the 8th International Conference on Computational Methods in Systems Biology*, CMSB '10, pages 55–65, New York, NY, USA, 2010. ACM. available online: [arXiv:1005.0747 \[q-bio.QM\]](#).
- [HRSS10] Monika Heiner, Christian Rohr, Martin Schwarick, and Stefan Streif. A comparative study of stochastic analysis techniques. In *Proceedings of the 8th International Conference on Computational Methods in Systems Biology*, CMSB '10, pages 96–106, New York, NY, USA, 2010. ACM.
- [KR81] John C. Kieffer and Maurice Rahe. Markov channels are asymptotically mean stationary. *SIAM J. Math. Anal.*, 12(3):293–305, 1981.
- [KS76] John G. Kemeny and J. Laurie Snell. *Finite Markov chains*. Springer-Verlag, New York, 1976. Reprinting of the 1960 original, Undergraduate Texts in Mathematics.
- [Par99] Emanuel Parzen. *Stochastic processes*, volume 24 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999. Reprint of the 1962 original.
- [Pin64] M. S. Pinsker. *Information and information stability of random variables and processes*. Translated and edited by Amiel Feinstein. Holden-Day Inc., San Francisco, Calif., 1964.
- [RP81] L. C. G. Rogers and J. W. Pitman. Markov functions. *Ann. Probab.*, 9(4):573–582, 1981.
- [Sar00] Ramesh R. Sarukkai. Link prediction and path analysis using Markov chains. *Computer Networks*, 33(1–6):377–386, 2000.
- [WA60] Satosi Watanabe and Chacko T. Abraham. Loss and recovery of information by coarse observation of stochastic chain. *Information and Control*, 3:248–278, 1960.
- [Wil11] D.J. Wilkinson. *Stochastic Modelling for Systems Biology*. Chapman & Hall/CRC Mathematical & Computational Biology. Taylor & Francis, 2011.
- [Woe09] Wolfgang Woess. *Denumerable Markov chains*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2009. Generating functions, boundary theory, random walks on trees.